arXiv:1412.0884v1 [math.GR] 2 Dec 2014

ON Ш-RIGIDITY OF GROUPS OF ORDER p^6

PRADEEP K. RAI AND MANOJ K. YADAV

Abstract. Let G be a group and $Out_c(G)$ be the group of its class-preserving outer automorphisms. We compute $|Out_c(G)|$ for all the group G of order p^6, where p is an odd prime. As an application, we observe that if G is a Ш-rigid group of order p^6, then it's Bogomolov multiplier $B_0(G)$ is zero.

1. Introduction

Let G be a group which acts on itself by conjugation, and let $H^1(G,G)$ be the first cohomology pointed set. Denote by $Ш(G)$ the subset of $H^1(G,G)$ consisting of the cohomology classes becoming trivial after restricting to every cyclic subgroup of G. The set $Ш(G)$, for a given group G, is called the Shafarevich-Tate set of G. Following Kunyavskiĭ [20], we say that G is a Ш-rigid group if the set $Ш(G)$ consists of one element. Throughout the paper Ш-rigid groups will be called rigid groups. The Bogomolov multiplier $B_0(G)$ of a finite group G is defined as the subgroup of the Schur multiplier consisting of the cohomology classes vanishing after restriction to all abelian subgroups of G. See [20] for further details on rigidity and Bogomolov multiplier of certain groups, and eventual ties between them. Kang and Kunyavskiĭ in [16] observed that $B_0(G) = 0$ for most of the known classes of finite Ш-rigid groups G, and asked the following question:

Question([16, Question 3.2]). Let G be a finite Ш-rigid group. Is it true that $B_0(G) = 0$?

This question has an affirmative answer for all p-groups of order at most p^5. For p-groups of order $\leq p^4$, it is proved in [16], and for groups of order p^5, $p > 3$, it follows from [22] and [31]. For $p = 2, 3$, it can be easily verified using GAP [9]. However this question does not have a positive solution in general (negative answer for a group of order 256 is provided in the paper [16] itself), it is interesting to explore for which classes of groups it has a positive solution. This paper is devoted to studying the rigidity property of groups of order p^6 for an odd prime p and answering the above question in affirmative for the class of groups considered here. Our viewpoint on this study is a bit different. We study rigidity problem through automorphisms of groups. We make it more precise here. For a finite group G, an automorphism of G is called (conjugacy) class-preserving if it maps each element of G to some conjugate of it. The set of all class-preserving automorphisms of G, denoted by $Aut_c(G)$, forms a normal subgroup of $Aut(G)$ and contains $Inn(G)$, the group of all inner automorphisms of G. Let $Out_c(G)$ denote the quotient group $Aut_c(G)/Inn(G)$. Then there is a bijection between $Out_c(G)$ and $Ш(G)$ for any finite group G [23, 2.12]. Therefore a finite group G is Ш-rigid if and only if $Out_c(G) = 1$. Both approaches, i.e., through the Shafarevich-Tate sets (cf. [17, 24, 25]) as well as $Out_c(G)$ (cf. [8, 12, 18, 19, 31]), have been explored by many mathematicians studying the rigidity of various classes of groups. As mentioned above, in the present paper we take the latter approach.

Groups of order p^6, p an odd prime, are classified in 43 isoclinism families by R. James [15], which are denoted by Φ_k for $1 \leq k \leq 43$. The concept of isoclinism was introduced by P. Hall [10]. That $Aut_c(G)$, for a non-abelian finite group G, is independent (upto isomorphism) of the choice of a group in a given isoclinism family is shown in [31, Theorem 4.1]. This result allows us to select and work with any group from each of 43 isoclinism families of groups of order p^6.

2010 *Mathematics Subject Classification.* 20D45, 20D15.

Key words and phrases. class-preserving automorphism, Ш-rigid group.

The following is the main result of this paper, which classifies rigid groups of order p^6, p being an odd prime, and size of $|Ш(G)| = |Out_c(G)|$ is computed for the groups G which are not rigid.

Theorem A. *Let G be a group of order p^6 for an odd prime p. Then $Out_c(G) \neq 1$ if and only if G belongs to one of the isoclinism families Φ_k for $k = 7, 10, 13, 15, 18, 20, 21, 24, 30, 36, 38, 39$. Moreover,*

1. *if G belongs to one of the isoclinism families Φ_k for $k = 7, 10, 24, 30, 36, 38, 39$, then $|Out_c(G)| = p$,*
2. *if G belongs to one of the isoclinism families Φ_k for $k = 13, 18, 20$, then $|Out_c(G)| = p^2$, and*
3. *if G belongs to one of the isoclinism families Φ_k for $k = 15, 21$, then $|Out_c(G)| = p^4$.*

Proof of Theorem A follows from Theorem 3.8 and Theorem 4.9. Using GAP [9] it can be verified that the above question has an affirmative answer for groups of order 2^6 and 3^6 (Bogomolov multiplier of groups of order 2^6 is also computed in [6]). With this information and results of Chen and Ma [5, Theorem 1.4], Theorem A provides an affirmative answer of the above mentioned question of Kang and Kunyavskiĭ for groups of order p^6 for all primes p in the following result.

Theorem B. *Let G be a Ш-rigid group of order p^6 for a prime p. Then its Bogomolov multiplier $B_0(G)$ is zero.*

Theorem A is also interesting from other point of view. In 1911 W. Burnside [3] asked the following question: Does there exist any finite group G such that G has a non-inner class-preserving automorphism? In 1913, he himself answered this question affirmatively by constructing a group W of order p^6, p an odd prime [4]. This group W is of nipotency class 2 such that $|Aut_c(W)| = p^8$ and $|Out_c(W)| = p^4$. More groups with non-inner class-preserving automorphism were constructed in [2, 11, 13, 21, 28]. We refer the reader to [30] for a more comprehensive survey on the topic. So Theorem A above classifies all groups of order p^6, p an odd prime, having non-inner class-preserving automorphisms.

We use the following notations. For a multiplicatively written group G let $x, y, a \in G$. Then $[x, y]$ denotes the commutator $x^{-1}y^{-1}xy$ and x^a denotes the conjugate of x by a i.e. $a^{-1}xa$. By $\langle x \rangle$ we denote the cyclic subgroup of G generated by x. By $Z(G)$ and $Z_2(G)$ we denote the center and the second center of G respectively. For $x \in G$, $C_G(x)$ denotes the centralizer of x in G. By $C_G(H)$ we denote the centralizer of H in G, where H is a subgroup of G. We write $\gamma_2(G)$ for the commutator subgroup of G. The group of all homomorphisms from a group H to an abelian group K is denoted by $Hom(H, K)$. For $x \in G$, x^G denotes the G-conjugacy class of x and $[x, G]$ denotes the set of all $[x, y]$, for $y \in G$. By C_p we denote the cyclic group of order p. The subgroup generated by all elements of order p in $Z(G)$ is denoted by $\Omega_1(Z(G))$.

2. Preliminaries

The following commutator identities are easy to prove.

Lemma 2.1. *Let G be a group and $x, y, z \in G$. Then*

1. *$[xy, z] = [x, z]^y [y, z]$, and*
2. *$[z, xy] = [z, y][z, x]^y$.*

An automorphism of a group G is called central if it acts trivially on the central quotient group $G/Z(G)$, or equivalently, if it commutes with all the inner automorphisms of G. The set of all central automorphisms of G forms a normal subgroup of $Aut(G)$. We denote this subgroup by $Autcent(G)$. The following lemma follows from [1].

Lemma 2.2. *Let G be a purely non-abelian finite p-group. Then $|Autcent(G)| = |Hom(G/\gamma_2(G), Z(G))|$.*

Lemma 2.3 ([31], Lemma 2.2)**.** *Let G be a finite p-group such that $Z(G) \leq [x, G]$ for all $x \in G - \gamma_2(G)$. Then $|Aut_c(G)| \geq |Autcent(G)||G/Z_2(G)|$.*

Lemma 2.4 ([13], Proposition 14.4)**.** *Let G be a finite group and H be an abelian normal subgroup of G such that G/H is cyclic. Then $Out_c(G) = 1$.*

Lemma 2.5 ([26], Lemma 5)**.** *For a group G, $C_{Aut_c(G)}(Inn(G)) = Z(Aut_c(G))$.*

Lemma 2.6 ([31], Lemma 2.6)**.** *Let G be a finite group. Let $x_1, x_2, \ldots, x_d$ be a minimal generating set for G. Then $|Aut_c(G)| \leq \prod_{i=1}^{d} |{x_i}^G|$.*

Lemma 2.7. *Let G be a purely non-abelian p-group, minimally generated by $\alpha_1, \alpha_2, \ldots, \alpha_t$. Suppose that $G/\gamma_2(G)$ is elementary abelian. If $\beta_i \in \Omega_1(Z(G))$ for $i = 1, \ldots, t$, then the map $\delta : \{\alpha_1, \alpha_2, \ldots, \alpha_t\} \to G$, defined as $\delta(\alpha_i) = \alpha_i\beta_i$, extends to a central automorphism of G.*

Proof. Note that the map $f_\delta : \{\alpha_i\gamma_2(G) |\ i = 1, \ldots, t\} \to \Omega_1(Z(G))$ defined as $\alpha_i\gamma_2(G) \to \beta_i$ for $i = 1, \ldots, t$, extends to a homomorphism from $G/\gamma_2(G)$ to $Z(G)$ because $G/\gamma_2(G)$ is elementary abelian. Now it follows from [1, Theorem 1] that the map $g \to gf_\delta(g)$ for all $g \in G$ is a central automorphism of G. This proves the lemma. □

Let G be a group minimally generated by $\alpha_1, \alpha_2, \ldots, \alpha_t$. Then note that any element of G can be written as $\eta\alpha_1^{k_1}\alpha_2^{k_2}\cdots\alpha_t^{k_t}$ for some $\eta \in \gamma_2(G)$ and some $k_1, k_2, \ldots, k_t \in \mathbb{Z}$. Also note that any conjugate of an α_i can be written as $\alpha_i[\alpha_i, \eta\alpha_1^{k_1}\alpha_2^{k_2}\cdots\alpha_t^{k_t}]$ for some $\eta \in \gamma_2(G)$ and some $k_1, k_2, \ldots, k_t \in \mathbb{Z}$. It follows that an automorphism δ of G is class-preserving if and only if for every $k_1, k_2, \ldots, k_t \in \mathbb{Z}$ and for all $\eta_1 \in \gamma_2(G)$, there exist $l_1, l_2, \ldots, l_t \in \mathbb{Z}$ and $\eta_2 \in \gamma_2(G)$ (depending on $k_1, k_2, \ldots, k_t$ and η_1) such that

$$\left[\eta_1\prod_{i=1}^{t}\alpha_i^{k_i},\ \ \eta_2\prod_{i=1}^{t}\alpha_i^{l_i}\right] = \left(\eta_1\prod_{i=1}^{t}\alpha_i^{k_i}\right)^{-1}\delta\left(\eta_1\prod_{i=1}^{t}\alpha_i^{k_i}\right).$$

We will be using these facts and Lemma 2.1 very frequently in the proofs without any further reference.

3. Groups G with trivial $Out_c(G)$

In this section, we deal with those groups G for which $Out_c(G)$ is trivial. The isoclinism family Φ_k of groups of order p^6 contains a group $\Phi_k(1^6)$ for certain values of k. These groups will be considered frequently throughout the paper.

Lemma 3.1. *Let G be the group $\Phi_{11}(1^6)$. Then $Out_c(G) = 1$.*

Proof. The group G is a special p-group, minimally generated by α_1, α_2 and α_3. The commutator subgroup $\gamma_2(G)$ is generated by $\beta_1 := [\alpha_2, \alpha_3], \beta_2 := [\alpha_3, \alpha_1], \beta_3 := [\alpha_1, \alpha_2]$. The conjugates of α_1, α_2 and α_3 are $\alpha_1\beta_3^t\beta_2^s$, $\alpha_2\beta_3^t\beta_1^r$ and $\alpha_3\beta_2^s\beta_1^r$ respectively, where r, s and t vary over $\mathbb{Z}$. Since the exponent of $\gamma_2(G)$ is p, it follows that $|\alpha_i^G| = p^2$ for $i = 1, 2, 3$. Therefore by Lemma 2.6, $|Aut_c(G)| \leq p^6$. Define a map $\delta : \{\alpha_1, \alpha_2, \alpha_3\} \to G$ such that $\alpha_1 \mapsto \alpha_1\beta_3^{t_1}\beta_2^{s_1}, \alpha_2 \mapsto \alpha_2\beta_3^{t_2}\beta_1^{r_2}$ and $\alpha_3 \mapsto \alpha_3\beta_2^{s_3}\beta_1^{r_3}$, for some $s_1, t_1, r_2, t_2, r_3, s_3 \in \mathbb{Z}$. By Lemma 2.7, this map extends to a central automorphism of G. Since δ fixes $\gamma_2(G)$ element-wise, for $k_1, l_1, m_1 \in \mathbb{Z}$ and $\eta \in \gamma_2(G)$,

$$\delta(\eta\alpha_1^{k_1}\alpha_2^{l_1}\alpha_3^{m_1}) = \eta\alpha_1^{k_1}\alpha_2^{l_1}\alpha_3^{m_1}\beta_1^{l_1r_2+m_1r_3}\beta_2^{k_1s_1+m_1s_3}\beta_3^{k_1t_1+l_1t_2}.$$

Therefore δ extends to a class-preserving automorphism if and only if for every $k_1, l_1, m_1 \in \mathbb{Z}$, and $\eta_1 \in \gamma_2(G)$ there exist k_2, l_2, m_2 (depending on k_1, l_1, m_1) and $\eta_2 \in \gamma_2(G)$ such that

$$[\eta_1\alpha_1^{k_1}\alpha_2^{l_1}\alpha_3^{m_1}, \eta_2\alpha_1^{k_2}\alpha_2^{l_2}\alpha_3^{m_2}] = \beta_1^{l_1r_2+m_1r_3}\beta_2^{k_1s_1+m_1s_3}\beta_3^{k_1t_1+l_1t_2}.$$

Expanding the left hand side, we get

$$\beta_1^{l_1m_2-l_2m_1}\beta_2^{k_2m_1-k_1m_2}\beta_3^{k_1l_2-k_2l_1} = \beta_1^{l_1r_2+m_1r_3}\beta_2^{k_1s_1+m_1s_3}\beta_3^{k_1t_1+l_1t_2}.$$

Comparing the powers of β_i's, we have that δ extends to a class-preserving automorphism if and only if the following equations hold true:

$$l_1m_2 - l_2m_1 \equiv l_1r_2 + m_1r_3 \pmod{p}$$

$$k_2m_1 - k_1m_2 \equiv k_1s_1 + m_1s_3 \pmod{p}$$

$$k_1l_2 - k_2l_1 \equiv k_1t_1 + l_1t_2 \pmod{p}.$$

Let δ be a class-preserving automorphism. Choose $k_1 = 0$ and m_1, l_1 to be non-zero modulo p. Then $k_2m_1 \equiv m_1s_3 \pmod{p}$ and $-k_2l_1 \equiv l_1t_2 \pmod{p}$. It follows that $t_2 \equiv -s_3 \pmod{p}$. Similarly if we choose l_1 to be zero and k_1, m_1 to be non-zero modulo p, then $r_3 \equiv -t_1 \pmod{p}$, and if m_1 to be zero and l_1, k_1 to be non-zero modulo p, then $r_2 \equiv -s_1 \pmod{p}$. It follows that $|Aut_c(G)| \leq p^3 = |Inn(G)|$. Therefore $Out_c(G) = 1$. □

Lemma 3.2. *Let G be one of the groups $\Phi_{17}(1^6)$ and $\Phi_{19}(1^6)$. Then $Out_c(G) = 1$.*

Proof. First assume that G is the group $\Phi_{17}(1^6)$. Then G is a p-group of nilpotency class 3, minimally generated by α, α_1 and β. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_2 := [\alpha_1, \alpha], \alpha_3 := [\alpha_2, \alpha]$ and $\gamma := [\beta, \alpha_1]$. The center $Z(G)$ is of order p^2, generated by α_3 and γ. It is easy to see that $|\alpha_1^G| \leq p^2, |\alpha^G| \leq p^2$ and $|\beta^G| = p$. Therefore, applying Lemma 2.6, we have $|Aut_c(G)| \leq p^5$. Define a map $\delta : \{\alpha, \alpha_1, \beta\} \to G$ such that $\alpha \mapsto \alpha, \alpha_1 \mapsto \alpha_1$ and $\beta \mapsto \alpha_1^{-1}\beta\alpha_1 = \beta\gamma$. Suppose that $|Aut_c(G)| = p^5$. Then δ must extend to a class-preserving automorphism of G. Hence there exist $\eta_1 (= \alpha_2^{r_1}\alpha_3^{s_1}\gamma^{t_1}$ say$) \in \gamma_2(G)$ and $k_1, l_1, m_1 \in \mathbb{Z}$ such that $[\alpha\beta,\ \eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}] = \gamma$. It is a routine calculation to show that

$$[\alpha\beta, \eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}] = \alpha_2^{-l_1}\alpha_3^{-r_1}\gamma^{l_1},$$

which can not be equal to γ for any value of l_1 and r_1. Hence δ can not be a class-preserving automorphism, and hence $|Aut_c(G)| \leq p^4$. But $|Inn(G)| = p^4$, so we have $Aut_c(G) = Inn(G)$.

Now we take G to be $\Phi_{19}(1^6)$. Then G is a p-group of class 3, minimally generated by α, α_1 and α_2. Define a map $\delta : \{\alpha, \alpha_1, \alpha_2\} \to G$ such that $\alpha \mapsto \alpha_1^{-1}\alpha\alpha_1 (= \alpha\beta_1), \alpha_1 \mapsto \alpha_1$ and $\alpha_2 \mapsto \alpha_2$. Now the proof follows on the same lines as for the group $\Phi_{17}(1^6)$. □

Lemma 3.3. *Let G be the group $\Phi_{23}(1^6)$. Then $Out_c(G) = 1$.*

Proof. The group G is a p-group of class 4, minimally generated by α, α_1. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_{i+1} := [\alpha_i, \alpha]$ for $1 \leq i \leq 3$ and $\gamma := [\alpha_1, \alpha_2]$. The center $Z(G)$ is of order p^2, generated by α_4 and γ. It is easy to check that $|\alpha^G| \leq p^3$ and $|\alpha_1^G| \leq p^2$. Therefore $|Aut_c(G| \leq p^5$. Let $H = \langle\alpha_4\rangle$. Since $\alpha_4 \in Z(G), H$ is normal. Consider the quotient group G/H. One can check that the group G/H belongs to the family Φ_6. Therefore it follows from [31, Lemma 5.3] that $Aut_c(G/H) = Inn(G/H)$. Now define a map $\delta : \{\alpha, \alpha_1\} \to G$, such that $\delta(\alpha) = \alpha$ and $\delta(\alpha_1) = \alpha_2^{-1}\alpha_1\alpha_2 = \alpha_1\gamma$. Suppose that $|Aut_c(G| = p^5$. Then δ must extend to a class-preserving automorphism of G. It also induces a non-trivial class-preserving automorphism (say $\bar{\delta}$) of G/H. But then $\bar{\delta}$ is an inner automorphism of G/H because $Aut_c(G/H) = Inn(G/H)$. Now note that $\bar{\delta}$ is also a central automorphism of G/H. It follows that it is induced by some element in $Z_2(G/H) = \langle\alpha_2H, Z(G/H)\rangle$. Let $\bar{\delta}$ be induced by α_2^tH for some $t \in \mathbb{Z}$. Hence

$$\bar{\delta}(\alpha H) = (\alpha_2^tH)^{-1}\alpha H(\alpha_2^tH) = \alpha\alpha_3^{-t}H.$$

This must be equal to αH. But then $t = 0 \pmod{p}$ and hence $\bar{\delta}(\alpha_1H) = \alpha_1H$, a contradiction. Therefore we have $|Aut_c(G)| \leq p^4$. But $|Inn(G)| = p^4$, therefore $Out_c(G) = 1$. □

Lemma 3.4. *Let G be the group $\Phi_{27}(1^6)$. Then $Out_c(G) = 1$.*

Proof. The group G is a p-group of class 4, minimally generated by α, α_1, β. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_{i+1} := [\alpha_i, \alpha]$ for $1 \leq i \leq 2$ and $\alpha_4 := [\alpha_3, \alpha] = [\alpha_1, \beta] = [\alpha_1, \alpha_2]$. The center $Z(G)$ is of order p, generated by α_4. It is easy to check that $|\alpha^G| \leq p^3, |\alpha_1^G| \leq p^2$ and $|\beta^G| = p$. Therefore $|Aut_c(G)| \leq p^6$. Define $\delta : \{\alpha, \alpha_1, \beta\} \to G$ such that $\delta(\alpha) = \alpha, \delta(\alpha_1) = \alpha_1$ and $\delta(\beta) = \alpha_1\beta\alpha_1^{-1} = \beta\alpha_4$. Suppose that $|Aut_c(G)| = p^6$. Then δ extends to a class-preserving automorphism of G. Also note that δ fixes α_2, therefore $\delta(\alpha_2\beta^{-1}) = \alpha_2\beta^{-1}\alpha_4^{-1}$. Since δ is a class-preserving automorphism, there exist some $g \in G$ such that $[\alpha_2\beta^{-1}, g] = \alpha_4^{-1}$. Note that $C_G(\alpha_2\beta^{-1}) = \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \beta \rangle$. Therefore without loss of generality we can assume that $g = \alpha^{k_1}$. It can be calculated that

$$[\alpha_2\beta^{-1}, \alpha^{k_1}] = \alpha_3^{k_1}\alpha_4^{k_1(k_1-1)/2},$$

which, for any value of k_1, can never be equal to α_4^{-1}. Hence, we get a contradiction. Therefore $|Aut_c(G)| \leq p^5$. But $|Inn(G)| = p^5$, therefore $Out_c(G) = 1$. □

Lemma 3.5. *Let $G \in \{\Phi_{28}(222), \Phi_{29}(222)\}$. Then $|Out_c(G)| = 1$.*

Proof. The group G is a p-group of class 4, minimally generated by α and α_1. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_{i+1} := [\alpha_i, \alpha]$ for $1 \leq i \leq 2$ and $\alpha_4 := [\alpha_3, \alpha] = [\alpha_1, \alpha_2]$. The center $Z(G)$ is of order p, generated by α_4. Note that $\langle \alpha, \alpha_4 \rangle \leq C_G(\alpha)$ and hence $|C_G(\alpha)| \geq p^3$, therefore $|\alpha^G| \leq p^3$. Now note that, since $\alpha_2^{(p)} = \alpha_4^y$, we have for $p = 3, \alpha_2^p = \alpha_4^{y-1}$, and for $p > 3, \alpha_2^p = \alpha_4^y$. Following is a routine calculation.

$$[\alpha_1, \alpha^p] = \alpha_2^p\alpha_3^{p(p-1)/2}\alpha_4^{\sum_{n=1}^{p-2} n(n-1)/2} = \alpha_2^p\alpha_4^{(p-2)(p-1)p/3},$$

which, for $p = 3$, equals α_4^{y+1}, and for $p > 3$, equals α_4^y. But we have $[\alpha_1, \alpha_2^t] = \alpha_4^t$, therefore $\alpha_2^{y+1}\alpha^{-p} \in C_G(\alpha_1)$ for $p = 3$ and $\alpha_2^y\alpha^{-p} \in C_G(\alpha_1)$ for $p > 3$. Now it is easy to see that $|C_G(\alpha_1)| \geq p^4$. Hence $|\alpha_1^G| \leq p^2$. It follows from Lemma 2.6 that $|Aut_c(G)| \leq p^5$. But $|G/Z(G)| = p^5$, therefore $Out_c(G) = 1$. □

Lemma 3.6. *Let $G \in \{\Phi_k(1^6), \Phi_{34}(321)a \mid \ k = 31, \ldots, 33\}$. Then $Out_c(G) = 1$.*

Proof. The group G is a p-group of class 3, minimally generated by α, α_1 and α_2. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\beta_i := [\alpha_i, \alpha]$ for $i = 1, 2$ and $\gamma := [\alpha_1, \beta_1]$. For $k = 31, 32$, $[\alpha_2, \beta_2] = \gamma^y$, and for $k = 33$ and $\Phi_{34}(321)a$, $\gamma = [\beta_2, \alpha]$. The center $Z(G)$ is of order p, generated by γ. It is easy to see that, if $G \in \{\Phi_k(1^6) \mid k = 31, 32\}$, then $|\alpha_1^G| \leq p^2, |\alpha_2^G| \leq p^2, |\alpha^G| \leq p^2$, and if $G \in \{\Phi_{33}(1^6), \Phi_{34}(321)a\}$, then $|\alpha_1^G| \leq p^2, |\alpha_2^G| = p$ and $|\alpha^G| \leq p^3$. Therefore for all the four groups G, $|Aut_c(G)| \leq p^6$. Define a map $\delta : \{\alpha, \alpha_1, \alpha_2\} \to G$ such that $\alpha \mapsto \alpha, \alpha_1 \mapsto \alpha_1$ and $\alpha_2 \mapsto \alpha^{-1}\alpha_2\alpha = \alpha_2\beta_2$. Suppose that $|Aut_c(G)| = p^6$. Then δ must extend to a class-preserving automorphism of G. Hence there exist elements $\eta_1 \in \gamma_2(G)$ and $k_1, l_1, m_1 \in \mathbb{Z}$ such that $[\alpha_1\alpha_2,\ \eta_1\alpha^{k_1}\alpha_1^{l_1}\alpha_2^{m_1}] = \beta_2$, but it is a routine calculation that

$$[\alpha_1\alpha_2,\ \eta_1\alpha^{k_1}\alpha_1^{l_1}\alpha_2^{m_1}] = \beta_1^{k_1}\beta_2^{k_1}\gamma^a$$

for some $a \in \mathbb{Z}$. Clearly it can not be equal to β_2 for any value of k_1, l_1, m_1, r_1 and s_1. Thus δ is not a class-preserving automorphism, and therefore it follows that $|Aut_c(G)| \leq p^5$. But $|Inn(G)| = p^5$. Hence $Aut_c(G) = Inn(G)$ proving that $Out_c(G) = 1$. □

Lemma 3.7. *Let $G \in \{\Phi_k(1^6), \Phi_j(222)a_0 \mid k = 40, 41,\ j = 42, 43\}$. Then $Out_c(G) = 1$.*

Proof. The group G is a p-group of class 4, minimally generated by α_1 and α_2. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\beta := [\alpha_1, \alpha_2], \beta_i := [\beta, \alpha_i]$ for $i = 1, 2$ and γ, where, for $k = 40$, $\gamma := [\beta_1, \alpha_2] = [\beta_2, \alpha_1]$, for $k = 41$, $\gamma^{-\nu} := [\alpha_2, \beta_2] = [\alpha_1, \beta_1]^{-\nu}$, for $j = 42$, $\gamma := [\alpha_1, \beta_2] = [\alpha_2, \beta_1]$ and for $j = 43$, $\gamma^{-\nu} := [\alpha_2, \beta_2] = [\alpha_1, \beta_1]^{-\nu}$. The center $Z(G)$ is of order p, generated by γ. It is easy to see that $|\alpha_1^G| \leq p^3$ and $|\alpha_2^G| \leq p^3$. Therefore

$|Aut_c(G)| \leq p^6$. Define a map $\delta : \{\alpha_1, \alpha_2\} \to G$ such that $\alpha_1 \mapsto \alpha_1$ and $\alpha_2 \mapsto \alpha_2\beta_2$. Suppose that $|Aut_c(G)| = p^6$. Then δ must extend to a class-preserving automorphism of G. Hence there exist $\eta_1 (= \beta^{r_1}\beta_1^{s_1}\beta_2^{t_1}\gamma^{u_1}$ say) $\in \gamma_2(G)$ and $k_1, l_1 \in \mathbb{Z}$ such that $[\alpha_1\alpha_2,\ \eta_1\alpha_1^{k_1}\alpha_2^{l_1}] = \beta_2$. It is a routine calculation to show that,

$$[\alpha_1\alpha_2,\ \eta_1\alpha_1^{k_1}\alpha_2^{l_1}] \quad = \quad \beta^{l_1-k_1}\beta_1^{-k_1(k_1-1)/2-r_1}\beta_2^{l_1(l_1+1)/2-k_1l_1-r_1}\gamma^a$$

for some $a \in \mathbb{Z}$. It is easy to see that if powers of β and β_1 in the above expression are 0 modulo p, then the power of β_2 is also 0 modulo p. It follows that δ is not a class-preserving automorphism, and therefore $|Aut_c(G)| \leq p^5$. But $|Inn(G)| = p^5$, therefore $Out_c(G) = 1$. □

We are now ready to prove the following theorem.

Theorem 3.8. *Let G be a group of order p^6 which belongs to one of the isoclinism family Φ_k for $k = 2, \ldots, 6, 8, 9, 11, 12, 14, 16, 17, 19, 23, 25, \ldots, 29, 31, \ldots, 35, 37, 40, \ldots, 43$, [15, Section 4.6]. Then $Out_c(G) = 1$.*

In the following proof, $\Phi_k(1^5)$ is the group of order p^5 from the isoclinism family (k) of [15, Section 4.5] and $\Phi_8(32)$ is the group of order p^5 from the isoclinism family (8) of [15, Section 4.5].

Proof. Note that $Aut_c(H \times K) \cong Aut_c(H) \times Aut_c(K)$, for any two groups H and K. Let $G \in \{\Phi_k(1^6),\ \Phi_8(321)a,\ \mid k = 2, \ldots, 6, 9\}$, then, since $\Phi_k(1^6) = \Phi_k(1^5) \times C_p$ and $\Phi_8(321)a = \Phi_8(32) \times C_p$, it follows from [31, Theorem 5.5] that $Out_c(G) = 1$. Since the group $\Phi_{12}(1^6)$ is a direct product of groups of order p^3, $Out_c(\Phi_{12}(1^6)) = 1$. The group $\Phi_{14}(1^6)$ is of nilpotency class 2 and $\gamma_2(\Phi_{14}(1^6))$ is cyclic, therefore from [31, Corollary 3.6], we have $Out_c(\Phi_{14}(1^6)) = 1$. Note that if $G \in \{\Phi_{16}(1^6), \Phi_k(222) \mid k = 25, 26\}$, then G is abelian by cyclic. Hence by Lemma 2.4 $Out_c(G) = 1$. If $G \in \{\Phi_k(1^6) \mid k = 22, 35, 37\}$, then by Lemma 2.6 it follows that $|Aut_c(G)| \leq p^5$. Since $|Inn(G)| = p^5$, $Out_c(G) = 1$. This, along with lemmas 3.1 - 3.7, completes the proof of Theorem 3.8. □

4. Groups G with non-trivial $Out_c(G)$

In this section we deal with those groups for which there exist a non-inner class-preserving automorphism.

Lemma 4.1. *Let G be the group $\Phi_{24}(1^6)$. Then $|Out_c(G)| = p$.*

Proof. The group G is a p-group of class 4, minimally generated by α, α_1 and β. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_{i+1} := [\alpha_i, \alpha]$ for $i = 1, 2$ and $\alpha_4 := [\alpha_3, \alpha] = [\alpha_1, \beta]$. The center $Z(G)$ is of order p, generated by α_4. We will show that for every element $g \in G - \gamma_2(G), Z(G) \leq [g, G]$. Let $g = \eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}$. Then

$$[g, \beta^{l_2}] = [\eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}, \beta^{l_2}] = \alpha_4^{l_1l_2}.$$

Therefore, if l_1 is non-zero modulo p, we have $Z(G) \leq [g, G]$. Let $l_1 \equiv 0 \pmod p$. Then, since $\alpha_1^p \in \gamma_2(G)$,

$$[g, \alpha_3^{k_2}] = [\eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}, \alpha_3^{k_2}] = \alpha_4^{k_1k_2}.$$

Therefore, if k_1 is non-zero modulo p, $Z(G) \leq [g, G]$. Now let $k_1 \equiv 0 \pmod p$. Then

$$[g, \alpha_1^{m_2}] = [\eta_1\beta^{m_1}, \alpha_1^{m_2}] = \alpha_4^{-m_1m_2}.$$

Therefore, if m_1 is non-zero modulo p, $Z(G) \leq [g, G]$. It follows that for every $g \in G - \gamma_2(G), Z(G) \leq [g, G]$. Hence by lemmas 2.2 and 2.3 we have,

$$|Aut_c(G)| \geq |Autcent(G)||G|/|Z_2(G)| = p^3p^6/p^3 = p^6.$$

But, $|\alpha^G| \leq p^3, |\alpha_1^G| \leq p^2$ and $|\beta^G| = p$. Therefore we have $|Aut_c(G)| \leq p^6$. Hence $|Aut_c(G)| = p^6$. Since $|G/Z(G)| = p^5$, it follows that $|Out_c(G)| = p$. □

Lemma 4.2. *Let G be the group $\Phi_{30}(1^6)$. Then $|Out_c(G)| = p$.*

Proof. The group G is a p-group of class 4, minimally generated by α, α_1 and β. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_2 := [\alpha_1, \alpha]$, $\alpha_3 := [\alpha_2, \alpha] = [\alpha_1, \beta]$ and $\alpha_4 := [\alpha_3, \alpha] = [\alpha_2, \beta]$. The center $Z(G)$ is of order p, generated by α_4. It is easy to see that $|\alpha^G| \leq p^3, |\alpha_1^G| \leq p^2$, and $|\beta^G| \leq p^2$. Therefore $|Aut_c(G)| \leq p^7$. Define a map $\delta : \{\alpha, \alpha_1, \beta\} \to G$ such that $\alpha \mapsto \alpha\alpha_4, \alpha_1 \mapsto \alpha_1$ and $\beta \mapsto \beta\alpha_4$. By Lemma 2.7, the map δ extends to a central automorphism of G. We will show that δ is also a non-inner class-preserving automorphism. Let $g = \eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}$. Then

$$[g, \alpha_3^{k_2}] = [\eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}, \alpha_3^{k_2}] = \alpha_4^{k_1k_2}.$$

Therefore, if k_1 is non-zero modulo p, we have $Z(G) \leq [g, G]$. Hence δ maps g to a conjugate of g. Now let $k_1 \equiv 0 \pmod p$, then

$$[g, \alpha_2^{m_2}] = [\eta_1\alpha_1^{l_1}\beta^{m_1}, \alpha_2^{m_2}] = \alpha_4^{-m_1m_2}.$$

It follows that, if m_1 is non-zero modulo p, δ maps g to a conjugate of g. Now let $m_1 \equiv 0 \pmod p$. But then, $\delta(\eta_1\alpha_1^{l_1}) = \eta_1\alpha_1^{l_1}$, because being a central automorphism δ fixes $\gamma_2(G)$ element-wise. We have shown that for every $g \in G$, $\delta(g)$ is a conjugate of g. Therefore δ is a class-preserving automorphism. Now suppose δ is an inner automorphism. Then being central, δ is induced by some element in $Z_2(G) = \langle\alpha_3, \alpha_4\rangle$. Since $\alpha_4 \in Z(G)$, we can assume that δ is induced by α_3^t. But then $\delta(\beta) = \beta$, a contradiction. Therefore δ is non-inner class-preserving automorphism. Since $|Inn(G)| = p^5$, it follows that $|Aut_c(G)| \geq p^6$. Now define a map $\sigma : \{\alpha, \alpha_1, \beta\} \to G$ such that $\alpha \mapsto \alpha_1^{-1}\alpha\alpha_1 = \alpha\alpha_2^{-1}, \alpha_1 \mapsto \alpha_1$ and $\beta \mapsto \beta$. Suppose $|Aut_c(G)| = p^7$. Then σ extends to a class preserving automorphism. But then, $1 = \delta([\alpha, \beta]) = [\alpha\alpha_2^{-1}, \beta] = \alpha_4^{-1}$ which is not possible. Therefore, we have $|Aut_c(G)| = p^6$. Since $|G/Z(G)| = p^5$, we have $|Out_c(G)| = p$. □

Lemma 4.3. *Let G be the group $\Phi_{36}(1^6)$. Then $|Out_c(G)| = p$.*

Proof. The group G is a p-group of maximal class, minimally generated by α and α_1. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_{i+1} := [\alpha_i, \alpha]$ for $i = 1, 2, 3$ and $\alpha_5 := [\alpha_4, \alpha] = [\alpha_1, \alpha_2]$. The center $Z(G)$ is of order p, generated by α_5. We will show that for every element $g \in G - \gamma_2(G), Z(G) \leq [g, G]$. Let $g = \eta_1\alpha^{k_1}\alpha_1^{l_1}$. Then

$$[g, \alpha_4^{k_2}] = [\eta_1\alpha^{k_1}\alpha_1^{l_1}, \alpha_4^{k_2}] = \alpha_4^{-k_1k_2}.$$

Therefore, if k_1 is non-zero modulo p, we have $Z(G) \leq [g, G]$. Let $k_1 \equiv 0 \pmod p$. Then

$$[g, \alpha_2^{l_2}] = [\eta_1\alpha_1^{l_1}, \alpha_2^{l_2}] = \alpha_5^{l_1l_2}.$$

Therefore, if l_1 is non-zero modulo p, $Z(G) \leq [g, G]$. Let $l_1 \equiv 0 \pmod p$, then we have $g \in \gamma_2(G)$ because $\alpha_1^p \in \gamma_2(G)$. It follows that, for every $g \in G - \gamma_2(G), Z(G) \leq [g, G]$. Hence from lemmas 2.2 and 2.3 we get

$$|Aut_c(G)| \geq |Autcent(G)||G|/|Z_2(G)| = p^2p^6/p^2 = p^6.$$

But it is easy to see that $|\alpha^G| \leq p^4$ and $|\alpha_1^G| \leq p^2$. Therefore $|Aut_c(G)| \leq p^6$. Hence $|Aut_c(G)| = p^6$. Since $|G/Z(G)| = p^5$, we have $|Out_c(G)| = p$. □

Lemma 4.4. *Let G be the group $\Phi_{38}(1^6)$. Then $|Out_c(G)| = p$.*

Proof. The group G is a p-group of maximal class, minimally generated by α, α_1. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_{i+1} := [\alpha_i, \alpha]$ for $i = 1, 2, 3$ and $\alpha_5 := [\alpha_4, \alpha] = [\alpha_1, \alpha_3]$. Also $[\alpha_1, \alpha_2] = \alpha_4\alpha_5^{-1}$. The center $Z(G)$ is generated by α_5. We show that for every element $g \in G - \gamma_2(G), Z(G) \leq [g, G]$. Let $g = \eta_1\alpha^{k_1}\alpha_1^{l_1}$. Then

$$[g, \alpha_4^{k_2}] = [\eta_1\alpha^{k_1}\alpha_1^{l_1}, \alpha_4^{k_2}] = \alpha_5^{-k_1k_2}.$$

Therefore, if k_1 is non-zero modulo p, $Z(G) \leq [g, G]$. Hence, let $k_1 \equiv 0 \pmod{p}$. Then

$$[g, \alpha_3^{l_2}] = [\eta_1 \alpha_1^{l_1}, \alpha_3^{l_2}] = \alpha_5^{l_1 l_2}.$$

Therefore, if l_1 is non-zero modulo p, we have $Z(G) \leq [g, G]$. If $l_1 \equiv 0 \pmod{p}$, then we have $g \in \gamma_2(G)$ because $\alpha_1^p \in \gamma_2(G)$. It follows that, for every $g \in G - \gamma_2(G)$, $Z(G) \leq [g, G]$. Therefore applying lemmas 2.2 and 2.3 we get

$$|Aut_c(G)| \geq |Autcent(G)||G|/|Z_2(G)| = p^2 p^6/p^2 = p^6.$$

It is easy to check that $|\alpha^G| \leq p^4$ and $|\alpha_1^G| \leq p^3$. Hence the upper bound for $|Aut_c(G)|$ is p^7. Suppose that $|Aut_c(G)| = p^7$. Then the map δ defined on the generating set $\{\alpha, \alpha_1\}$ as $\delta(\alpha) = \alpha$ and $\delta(\alpha_1) = \alpha_2^{-1}\alpha_1\alpha_2 = \alpha_1\alpha_4\alpha_5^{-1}$ must extend to a class-preserving automorphism automorphism of G. Note that

$$\delta(\alpha_2) = \delta([\alpha_1, \alpha]) = [\alpha_1\alpha_4\alpha_5^{-1}, \alpha] = \alpha_2\alpha_5.$$

Since δ is a class-preserving automorphism, there exists an $\eta_1 \in \gamma_2(G)$ and $k_1, l_1 \in \mathbb{Z}$ such that

$$[\alpha_2, \eta_1 \alpha^{k_1} \alpha_1^{l_1}] = \alpha_5.$$

We have that

$$[\alpha_2, \eta_1 \alpha^{k_1} \alpha_1^{l_1}] = \alpha_3^{k_1} \alpha_4^{k_1(k_1-1)/2 - l_1} \alpha_5^{l_1 - k_1 l_1 + \sum_{n=1}^{k_1-2} n(n-1)/2},$$

which for no values of k and k_1 can be equal to α_5. This gives a contradiction. Hence $|Aut_c(G)| = p^6$. Since $|G/Z(G)| = p^5$, we have $|Out_c(G)| = p$. □

Lemma 4.5. *Let G be the group $\Phi_{39}(1^6)$. Then $|Out_c(G)| = p$.*

Proof. The group G is a p-group of maximal class, minimally generated by α, α_1. The commutator subgroup $\gamma_2(G)$ is generated by $\alpha_{i+1} := [\alpha_i, \alpha]$ for $i = 1, 2$, $\alpha_4 := [\alpha_3, \alpha] = [\alpha_1, \alpha_2]$, and $\alpha_5 := [\alpha_2, \alpha_3] = [\alpha_3, \alpha_1] = [\alpha_4, \alpha_1]$. Note that any element of $\gamma_2(G)$ can be written as $\alpha_2^r \alpha_3^s \alpha_4^t \alpha_5^u$ for some $r, s, t, u \in \mathbb{Z}$. The center $Z(G)$ is generated by α_5. It is easy to see that $|\alpha^G| \leq p^3$ and $|\alpha_1^G| \leq p^3$. Therefore $|Aut_c(G)| \leq p^6$. Define a map $\delta : \{\alpha, \alpha_1\} \to G$ such that $\alpha \mapsto \alpha$, and $\alpha_1 \mapsto \alpha_2^{-1}\alpha_1\alpha_2 = \alpha_1\alpha_4$. We will show that δ extends to a class preserving automorphism of G. It is easy to check that δ preserves all the defining relations of the group. Hence δ extends to an endomorphism. Note that δ fixes $\gamma_2(G)$ element-wise, therefore for $k_1, l_1 \in \mathbb{Z}$ and $\eta_1 = \alpha_2^{r_1}\alpha_3^{s_1}\alpha_4^{t_1}\alpha_5^{u_1} \in \gamma_2(G)$,

$$\delta(\eta_1 \alpha^{k_1} \alpha_1^{l_1}) = \eta_1 \alpha^{k_1} \alpha_1^{l_1} \alpha_4^{l_1} \alpha_5^{l_1(l_1-1)/2}.$$

It is a routine calculation that

$$[\eta_1 \alpha^{k_1} \alpha_1^{l_1}, \alpha_3^{s_2} \alpha_4^{t_2}] = \alpha_5^{r_1 s_2 - l_1 s_2 - l_1 t_2 - l_1 k_1 s_2} \alpha_4^{-k_1 s_2}.$$

Since $\delta(\eta_1 \alpha^{k_1}) = \eta_1 \alpha^{k_1}$, which is obviously a conjugate of $\eta_1 \alpha^{k_1}$, let l_1 be non-zero modulo p. Now if k_1 is non-zero modulo p, then clearly there exist s_2 and t_2 such that

$$-k_1 s_2 \equiv l_1 \pmod{p}$$

and

$$r_1 s_2 - l_1 s_2 - l_1 t_2 - l_1 k_1 s_2 \equiv l_1(l_1 - 1)/2 \pmod{p}.$$

Suppose $k_1 \equiv 0 \pmod{p}$, then it can be calculated that

$$[\eta_1 \alpha_1^{l_1}, \alpha_2^{r_2} \alpha_4^{t_2}] = \alpha_5^{-s_1 r_2 + r_2 l_1 (l_1-1)/2 - l_1 t_2} \alpha_4^{l_1 r_2}.$$

Since l_1 is non-zero modulo p, clearly there exist r_2 and t_2 such that

$$l_1 r_2 \equiv l_1 \pmod{p}$$

and

$$-s_1 r_2 + r_2 l_1 (l_1 - 1)/2 - l_1 t_2 \equiv l_1(l_1 - 1)/2 \pmod{p}.$$

It follows that δ maps every element of G to a conjugate of itself. Therefore δ is a bijection, and hence a class-preserving automorphism of G. Now we show that δ is a non-inner automorphism.

On the contrary, suppose that δ is an inner automorphism. Note that $g^{-1}\delta(g) \in Z_2(G)$. Thus it follows that δ is induced by some element in $Z_3(G) = \langle \alpha_3, \alpha_4, Z(G)\rangle$, where $Z_3(G)$ denotes the third term in the upper central series of G. Let it be induced by $\alpha_3^{s_1}\alpha_4^{t_1}$. Since $\delta(\alpha) = \alpha$, we have $\alpha_3^{-s_1}\alpha\alpha_3^{s_1} = \alpha$, which implies that $s_1 \equiv 0 \pmod p$. But then $\delta(\alpha_1) = \alpha_4^{-t_1}\alpha_1\alpha_4^{t_1} = \alpha_1\alpha_5^{-t_1}$, a contradiction. Therefore δ is a non-inner class preserving automorphism. Since $|Inn(G)| = p^5$, we have $|Aut_c G)| = p^6$, and therefore $|Out_c(G)| = p$. □

Lemma 4.6. *Let G be the group $\Phi_{13}(1^6)$. Then $|Out_c(G)| = p^2$.*

Proof. The group G is a special p-group, minimally generated by $\alpha_1, \alpha_2, \alpha_3$ and α_4. The commutator subgroup $\gamma_2(G)$ is generated by $\beta_1 := [\alpha_1, \alpha_2]$ and $\beta_2 := [\alpha_1, \alpha_3] = [\alpha_2, \alpha_4]$. The conjugates of $\alpha_1, \alpha_2, \alpha_3$ and α_4 are , $\alpha_1\beta_1^r\beta_2^s$, $\alpha_2\beta_1^r\beta_2^s, \alpha_3\beta_2^s$ and $\alpha_4\beta_2^s$ respectively where r and s varies over $\mathbb{Z}$. Since the exponent of $\gamma_2(G)$ is p, it follows that $|\alpha_1^G| = |\alpha_2^G| = p^2$ and $|\alpha_3^G| = |\alpha_4^G| = p$. Therefore by Lemma 2.6, $|Aut_c(G)| \leq p^6$.

Define a map $\delta : \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\} \to G$ such that $\alpha_1 \mapsto \alpha_1\beta_1^{r_1}\beta_2^{s_1}, \alpha_2 \mapsto \alpha_2\beta_1^{r_2}\beta_2^{s_2}$, $\alpha_3 \mapsto \alpha_3\beta_2^{s_3}$ and $\alpha_4 \mapsto \alpha_4\beta_2^{s_4}$, for some $r_1, s_1, r_2, s_2, s_3, s_4 \in \mathbb{Z}$. By Lemma 2.7 this map extends to a central automorphism of G. Since δ fixes $\gamma_2(G)$ element-wise, for $k_1, l_1, m_1, n_1 \in \mathbb{Z}$ and $\eta_1 \in \gamma_2(G)$,

$$\delta(\eta_1\alpha_1^{k_1}\alpha_2^{l_1}\alpha_3^{m_1}\alpha_4^{n_1}) = \eta\alpha_1^{k_1}\alpha_2^{l_1}\alpha_3^{m_1}\alpha_4^{n_1}\beta_1^{k_1r_1+l_1r_2}\beta_2^{k_1s_1+l_1s_2+m_1s_3+n_1s_4}.$$

Therefore δ extends to a class-preserving automorphism if and only if for every $k_1, l_1, m_1, n_1 \in \mathbb{Z}$, and $\eta_1 \in \gamma_2(G)$, there exist k_2, l_2, m_2, n_2 (depending on k_1, l_1, m_1, n_1) and $\eta_2 \in \gamma_2(G)$ such that

$$[\eta_1\alpha_1^{k_1}\alpha_2^{l_1}\alpha_3^{m_1}\alpha_4^{n_1}, \eta_2\alpha_1^{k_2}\alpha_2^{l_2}\alpha_3^{m_2}\alpha_4^{n_2}] = \beta_1^{k_1r_1+l_1r_2}\beta_2^{k_1s_1+l_1s_2+m_1s_3+n_1s_4}.$$

Expanding the left hand side, we get

$$\beta_1^{k_1l_2-k_2l_1}\beta_2^{k_1m_2-k_2m_1+l_1n_2-l_2n_1} = \beta_1^{k_1r_1+l_1r_2}\beta_2^{k_1s_1+l_1s_2+m_1s_3+n_1s_4}.$$

Comparing the powers of β_i's, we see that δ extends to a class-preserving automorphism if the following equations hold:

$$k_1l_2 - k_2l_1 \equiv k_1r_1 + l_1r_2 \pmod p,$$
$$k_1m_2 - k_2m_1 + l_1n_2 - l_2n_1 \equiv k_1s_1 + l_1s_2 + m_1s_3 + n_1s_4 \pmod p.$$

It is easy to see that for any given k_1, l_1, m_1, n_1 there exist k_2, l_2, m_2, n_2 such that the above two equations are satisfied. Thus it follows that $|Aut_c(G)| = p^6$. Since $|G/Z(G)| = p^4, |Out_c(G)| = p^2$. □

Lemma 4.7. *Let G be the group $\Phi_{18}(1^6)$. Then $|Out_c(G)| = p^2$.*

Proof. The group G is a p-group of class 3, minimally generated by α, α_1, β. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\alpha_2 := [\alpha_1, \alpha], \alpha_3 := [\alpha_2, \alpha] = [\alpha_1, \beta]$ and $\gamma := [\alpha, \beta]$. The center $Z(G)$ is of order p^2, generated by α_3 and γ. Note that $|\alpha^G| \leq p^3, |\alpha_1^G| \leq p^2$ and $|\beta^G| \leq p^2$. It follows from Lemma 2.6 that $|Aut_c(G)| \leq p^7$. Now define a map $\delta : \{\alpha, \alpha_1, \beta\} \to G$ such that $\alpha \mapsto \alpha\alpha_3^{r_1}\gamma^{s_1}, \alpha_1 \mapsto \alpha_1\alpha_3^{r_2}$ and $\beta \mapsto \beta\alpha_3^{r_3}$ for some $r_1, s_1, r_2, r_3 \in \mathbb{Z}$. By Lemma 2.7 this map extends to a central automorphism of G. Let $g = \eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}$, where $\eta_1 \in \gamma_2(G)$ and $k_1, l_1, m_1 \in \mathbb{Z}$. Let $k_1 \equiv 0 \pmod p$. Then note that $\delta(g) = g\alpha_3^r$ for some $r \in \mathbb{Z}$ and

$$[g, \beta^{l_2}] = [\eta_1\alpha_1^{l_1}\beta^{m_1}, \beta^{l_2}] = \alpha_3^{l_1l_2}.$$

Therefore if l_1 is non-zero modulo p, we have $\langle\alpha_3\rangle \leq [g, G]$. Let $l_1 \equiv 0 \pmod p$. Note that $\alpha_1^p \in Z(G)$, hence

$$[g, \alpha_1^{m_2}] = [\eta_1\beta^{m_1}, \alpha_1^{m_2}] = \alpha_3^{-m_1m_2}.$$

Therefore if m_1 is non-zero modulo p, we have $\langle\alpha_3\rangle \leq [g, G]$. Thus we have shown that, if $k_1 \equiv 0 \pmod p$, then $g^{-1}\delta(g) \in [g, G]$. It follows that, if $k_1 \equiv 0 \pmod p$, δ maps g to a conjugate of g. Now suppose k_1 is non-zero modulo p. Then

$$[g, \alpha_2^{k_2}] = [\eta_1\alpha^{k_1}\alpha_1^{l_1}\beta^{m_1}, \alpha_2^{k_2}] = \alpha_3^{-k_1k_2},$$

so that $\langle \alpha_3 \rangle \leq [g, Z_2(G)]$. Also, we have

$$[g, \beta^{n_2}] = [\eta_1 \alpha^{k_1} \alpha_1^{l_1} \beta^{m_1}, \beta^{n_2}] = \alpha_3^{l_1 n_2} \gamma^{k_1 n_2}.$$

Since $\beta \in Z_2(G)$ and $[g, Z_2(G)]$ is a subgroup, it follows that $Z(G) \leq [g, G]$. Because δ is a central automorphism, we have $g^{-1}\delta(g) \in [g, G]$. We have shown that, for every $g \in G$, $g^{-1}\delta(g) \in [g, G]$. It follows that δ is a class-preserving automorphism. Since r_1, s_1, r_2, r_3 were arbitrary, we have that $|Aut_c(G) \cap Autcent(G)| \geq p^4$. Applying Lemma 2.5 we get $|Z(Aut_c(G))| \geq p^4$. Note that $Aut_c(G)$ is non-abelian because G is a group of class 3. Therefore $|Aut_c(G)| \geq p^6$. Now suppose that $|Aut_c(G)| = p^7$. Then the map σ defined on the generating set $\{\alpha, \alpha_1, \beta\}$ as $\alpha \mapsto \alpha, \alpha_1 \mapsto \alpha_1$ and $\beta \mapsto \alpha\beta\alpha^{-1} = \beta\gamma$ extends to a class-preserving automorphism. Hence there exist $\eta_2 \in \gamma_2(G)$ and $k_2, l_2, m_2 \in \mathbb{Z}$ such that $[\alpha_1\beta,\ \eta_2\alpha^{k_2}\alpha_1^{l_2}\beta^{m_2}] = \gamma$, but by a routine calculation it can be checked that

$$[\alpha_1\beta,\ \eta_2\alpha^{k_2}\alpha_1^{l_2}\beta^{m_2}] = \alpha_2^{k_2}\beta\alpha_3^{m_2 - l_2 + k_2(k_2-1)/2}\gamma^{-k_2},$$

which can not be equal to γ for any values of k_2, l_2 and m_2. Therefore we get a contradiction and it follows that $|Aut_c(G)| = p^6$. Hence $|Out_c(G)| = p^2$ as $|Inn(G)| = p^4$. □

Lemma 4.8. *Let G be the group $\Phi_{20}(1^6)$. Then $|Out_c(G)| = p^2$.*

Proof. The group G is a p-group of class 3, minimally generated by $\alpha, \alpha_1, \alpha_2$. The commutator subgroup $\gamma_2(G)$ is abelian and generated by $\beta := [\alpha_1, \alpha_2], \beta_1 := [\beta, \alpha_1]$ and $\beta_2 := [\beta, \alpha_2] = [\alpha, \alpha_1]$. The center $Z(G)$ is of order p^2, generated by β_1 and β_2. Note that $|\alpha^G| = p, |\alpha_1^G| \leq p^3$ and $|\alpha_2^G| \leq p^2$. It follows from Lemma 2.6 that $|Aut_c(G)| \leq p^6$. Now define a map $\delta : \{\alpha, \alpha_1, \alpha_2\} \to G$ such that $\alpha \mapsto \alpha\beta_2^{t_1}, \alpha_1 \mapsto \alpha_1\beta_1^{s_2}\beta_2^{t_2}$ and $\alpha_2 \mapsto \alpha_2\beta_2^{t_3}$ for some $t_1, s_2, t_2, t_3 \in \mathbb{Z}$. By Lemma 2.7 this map extends to a central automorphism of G. Let $g = \eta_1\alpha^{k_1}\alpha_1^{l_1}\alpha_2^{m_1}$, where $\eta_1 = \beta^{u_1}\beta_1^{v_1}\beta_2^{w_1}$ and $k_1, l_1, m_1, u_1, v_1, w_1 \in \mathbb{Z}$. Note that, if $l_1 \equiv 0 \pmod p$, then $\delta(g) = g\beta_2^r$ for some $r \in \mathbb{Z}$. Consider

$$[g, \beta^{m_2}] = [\eta_1\alpha^{k_1}\alpha_2^{m_1}, \beta^{m_2}] = \beta_2^{-m_1 m_2}.$$

Therefore if m_1 is non-zero modulo p, we have $\langle \beta_2 \rangle \leq [g, G]$. Let $m_1 \equiv 0 \pmod p$. Note that $\alpha_2^p \in Z(G)$, hence

$$[g, \alpha_2^{u_2}] = [\eta_1\alpha^{k_1}, \alpha_2^{u_2}] = [\beta^{u_1}\beta_1^{v_1}\beta_2^{w_1}\alpha^{k_1},\ \alpha_2^{u_2}] = \beta_2^{u_1 u_2}.$$

Therefore if u_1 is non-zero modulo p, we have $\langle \beta_2 \rangle \leq [g, G]$. Let $u_1 \equiv 0 \pmod p$. Then

$$[g, \alpha_1^{k_2}] = [\alpha^{k_1}, \alpha_1^{k_2}] = \beta_2^{k_1 k_2},$$

so that if k_1 is non-zero modulo p, then $\langle \beta \rangle \leq [g, G]$. Thus we have shown that if $l_1 \equiv 0 \pmod p$, then $g^{-1}\delta(g) \in [g, G]$. It follows that if $l_1 \equiv 0 \pmod p$, then δ maps g to a conjugate of g. Now suppose that l_1 is non-zero modulo p. Then

$$[g, \alpha^{l_2}] = [\eta_1\alpha^{k_1}\alpha_1^{l_1}\alpha_2^{m_1}, \alpha^{l_2}] = \beta_2^{-l_1 l_2},$$

so that $\langle \beta_2 \rangle \leq [g, Z_2(G)]$. Also we have

$$[g, \beta^{n_2}] = [\eta_1\alpha^{k_1}\alpha_1^{l_1}\alpha_2^{m_1}, \beta^{n_2}] = \beta_1^{-l_1 n_2}\beta_2^{-m_1 n_2}.$$

Since $\beta \in Z_2(G)$ and $[g, Z_2(G)]$ is a subgroup, it follows that $Z(G) \leq [g, G]$. Because δ is a central automorphism, we have $g^{-1}\delta(g) \in [g, G]$. It follows that δ is a class-preserving automorphism of G. Since t_1, s_2, t_2, t_3 were arbitrary, we have that $|Aut_c(G) \cap Autcent(G)| \geq p^4$. Applying Lemma 2.5 we get $|Z(Aut_c(G))| \geq p^4$. But $Aut_c(G)$ is non-abelian since G is of class 3. Therefore $|Aut_c(G)| \geq p^6$. Hence $|Aut_c(G)| = p^6$. Since $|G/Z(G)| = p^4$, we have $|Out_c(G)| = p^2$. □

Now we are ready to prove the following result, which together with Theorem 3.8 proves Theorem A.

Theorem 4.9. *Let G be a group of order p^6. Then the following hold true.*

(1) *If G belongs to any of the isoclinism families Φ_k for $k = 7, 10, 24, 30, 36, 38, 39$, then $|Out_c(G)| = p$.*

(2) *If G belongs to any of the isoclinism families Φ_k for $k = 13, 18, 20$, then $|Out_c(G)| = p^2$.*
(3) *If G belongs to any of the isoclinism families Φ_k for $k = 15, 21$, then $|Out_c(G)| = p^4$.*

Proof. Let G be either the group $\Phi_7(1^6)$ or the group $\Phi_{10}(1^6)$. Then, since $\Phi_7(1^6) = \Phi_7(1^5) \times C_p$ and $\Phi_{10}(1^6) = \Phi_{10}(1^5) \times C_p$, it follows from [31, Lemma 5.1 and Lemma 5.2] that $|Out_c(G)| = p$. With this observation in hands, (i) follows from lemmas 4.1-4.5. It is readily seen that (ii) follows from lemmas 4.6-4.8. Now let G be the group $\Phi_{15}(1^6)$. We observe that James' list of groups of order p^6 and class 2 consists of exactly 5 isoclinism families Φ_k for $k = 11, \ldots, 15$. As we have seen above that $|Aut_c(H)| \neq p^8$ for $H \in \{\Phi_k \mid 11 \leq k \leq 14\}$. But, as shown by Burnside [4], there exists a group W of order p^6 and of nilpotency class 2 such that $|Aut_c(W)| = p^8$. Therefore G must be isoclinic to the group W and $|Aut_c(G)| = p^8$. Since $|G/Z(G)| = p^4$, we have $|Out_c(G)| = p^4$. Next, it follows from [29, Proposition 5.8], that $|Out_c(\Phi_{21}(1^6))| = p^4$. This completes the proof of the theorem. □

Acknowledgements. We thank the referee for his/her useful comments and suggestions which make the paper more readable.

References

[1] J. E. Adney and T. Yen, *Automorphisms of a p-group*, Illinois J. Math **9** (1965), 137-143.
[2] P. A. Brooksbank and M.S. Mizuhara, *On groups with a class preserving outer automorphism.* Involve, **7** (2013), No. 2, 171-179.
[3] W. Burnside, *Theory of groups of finite order*, 2nd Ed. Dover Publications, Inc., 1955. Reprint of the 2nd edition (Cambridge, 1911).
[4] W. Burnside, *On the outer automorphisms of a p-group*, Proc. London Math. Soc. (2) **11** (1913), 40-42.
[5] Y.Chen and R. Ma, *Bogomolov multipliers of some groups of order p^6*, (Available at arXiv:1302.0584v1).
[6] H. Chu, S. Hu, M. Kang and B. Kunyavskiĭ, *Noether's problem and the unramified Brauer groups for groups of order 64*, Intern. Math. Res. Notices **12** (2010), 2329-2366.
[7] G. Ellis, *GAP package HAP - Homological Algebra Programming, Version 1.10.15*; 2014, (http://www.gap-system.org/Packages/hap.html).
[8] W. Feit and G. M. Seitz, *On finite rational groups and related topics*, Illinois J. Math. **33** (1989), no. 1, 103-131.
[9] The GAP Group, GAP - *Groups, Algorithms, and Programming, Version 4.7.4*; 2014, (http://www.gap-system.org).
[10] P. Hall, *The classification of prime power groups*, J. Reine Angew. Math. **182** (1940), 130-141.
[11] H. Heineken, *Nilpotente Gruppen, deren sämtliche Normalteiler charakteristisch sind*, Arch. Math. (Basel) **33** (1980), No. 6, 497-503.
[12] A. Herman and Y. Li, *Class preserving automorphisms of Blackburn groups*, J. Aust. Math. Soc. **80** (2006), 351-358.
[13] M. Hertweck, *Contributions to the integral representation theory of groups*, Habilitationsschrift, University of Stuttgart (2004). Available at http://elib.uni-stuttgart.de/opus/volltexte/2004/1638.
[14] M. Hertweck, *Class-preserving automorphisms of finite groups*, J. Algebra **241** (2001), 1-26.
[15] R. James, *The groups of order p^6 (p an odd prime)*, Math. Comp. **34** (1980), 613-637.
[16] M. Kang and B. Kunyavskiĭ, *The Bogomolov multiplier of rigid finite groups*, Arch. Math. (Basel) **102** (2014), 209-218.
[17] M. Kumar and L. R. Vermani, *"Hasse principle" for extraspecial p-groups*, Proc. Japan Acad. Ser. A **76** (2000), 123-125.
[18] M. Kumar and L. R. Vermani, *"Hasse principle" for groups of order p^4*, Proc. Japan Acad. Ser. A **77** (2001), 95-98.
[19] M. Kumar and L. R. Vermani, *On automorphisms of some p-groups*, Proc. Japan Acad. Ser. A **78** (2002), 46-50.
[20] B. Kunyavskiĭ, *Local-global invariants of finite and infinite groups: Around Burnside from another side*, Expo. Math. **31** (2013), 256-273.
[21] I. Malinowska, *On quasi-inner automorphisms of a finite p-group*, Publ. Math. Debrecen **41** (1992), No. 1-2, 73-77.
[22] P. Moravec, *Groups of order p^5 and their unramified Brauer groups*, J. Algebra **372** (2012), 320-327.
[23] T. Ono, *"Shafarevich-Tate sets" for profinite groups*, Proc. Japan Acad. Ser. A **75** (1999), 96-97.
[24] T. Ono and H. Wada, *"Hasse principle" for free groups*, Proc. Japan Acad. Ser. A **75** (1999), 1-2.
[25] T. Ono and H. Wada, *"Hasse principle" for symmetric and alternating groups*, Proc. Japan Acad. Ser. A **75** (1999), 61-62.
[26] P. K. Rai, *On class-preserving automorphisms of groups*, Ricerche mat. **63** (2014), 189-194.

[27] C. H. Sah, *Automorphisms of finite groups*, Journal of Algebra **10** (1968), 47-68.
[28] F. Szechtman, *n-inner automorphisms of finite groups*, Proc. Amer. Math. Soc. **131** (2003), 3657-3664.
[29] M. K. Yadav, *Class preserving automorphisms of finite p-groups*, J. London Math. Soc. **75(3)** (2007), 755-772.
[30] M. K. Yadav, *Class preserving automorphisms of finite p-groups - A Survey*, Proc. Groups St. Andrews (Bath) 2009, LMS Lecture Note Series **388** (2011), 569 - 579.
[31] M. K. Yadav, *On automorphisms of some finite p-groups*, Proc. Indian Acad. Sci. (Math. Sci.) **118(1)** (2008), 1-11.

School of Mathematics, Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Allahabad 211019, INDIA.
E-mail address: pradeeprai@hri.res.in

School of Mathematics, Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Allahabad 211019, INDIA.
E-mail address: myadav@hri.res.in